

**Editorial****Dual Inhibition of Cyclooxygenase & Lipoxygenase by Natural Products****Abdul Jabbar Shah**

Department of Pharmacy, COMSATS University Islamabad, Abbottabad Campus, Pakistan

\*Correspondence: [jabbarshah@cuiatd.edu.pk](mailto:jabbarshah@cuiatd.edu.pk)

© The Author(s) 2023. This article is licensed under a Creative Commons Attribution 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Cyclooxygenase (COX) and lipoxygenase (LOX) are enzymes involved in the metabolism of arachidonic acid, a fatty acid present in cell membranes. COX enzymes convert arachidonic acid into prostaglandins and thromboxanes, which are involved in inflammation, pain, and other physiological processes. On the other hand, LOX enzymes convert arachidonic acid into leukotrienes, which also play a role in inflammation and immune responses.

Dual inhibition of COX and LOX pathways has therapeutic implications, especially in conditions characterized by inflammation. Several natural products have been found to possess the ability to inhibit both COX and LOX enzymes simultaneously.

Curcumin, a compound found in turmeric, has been shown to exhibit dual inhibitory effects on COX and LOX enzymes. It can suppress the activity of both COX-1 and COX-2 enzymes, reducing the production of prostaglandins. Additionally, curcumin inhibits LOX enzymes, thereby reducing the synthesis of leukotrienes. This dual inhibition contributes to the anti-inflammatory effects of curcumin.

Resveratrol is a polyphenol compound found in various plant sources, including grapes and berries. It has been reported to possess COX and LOX inhibitory activity. Resveratrol inhibits COX enzymes, leading to a decrease in prostaglandin production. It also inhibits the activity of LOX enzymes, reducing leukotriene synthesis. The

anti-inflammatory properties of resveratrol have been studied in various inflammatory conditions. Green tea catechins: Green tea catechins, such as epigallocatechin gallate (EGCG), are bioactive compounds present in green tea. EGCG has been shown to inhibit both COX and LOX enzymes. It inhibits COX-2 expression and suppresses prostaglandin production. Moreover, EGCG inhibits the activity of LOX enzymes, reducing the synthesis of leukotrienes. Green tea catechins have demonstrated anti-inflammatory effects in preclinical and clinical studies.

Gingerol is a bioactive compound found in ginger. It has been reported to possess dual inhibitory effects on COX and LOX enzymes. Gingerol inhibits COX-2 activity and decreases prostaglandin production. It also inhibits LOX enzymes, reducing the synthesis of leukotrienes. Gingerol's anti-inflammatory properties have been studied in various models of inflammation.

It is important to note that the effectiveness and specificity of natural products in inhibiting COX and LOX enzymes may vary. The bioavailability and metabolism of these compounds can also influence their overall impact. Further research is necessary to fully understand the mechanisms and therapeutic potential of natural products in dual COX-LOX inhibition.

**Professor Dr. Abdul Jabbar Shah***Editor-In-Chief***Phytopharmacological Communication**