

Sensory Evaluation of Rice Grain: Methods, Applications, and Significance in Experimental Research

Farah Shamim*¹, Mohsin Ali Raza¹, Muhammad Shahid Munir¹

¹Rice Research Institute, Kala Shah Kaku, 39018, Lahore, Pakistan

Correspondence: farah_tirmazi@yahoo.com

ABSTRACT

Sensory evaluation is a dynamic tool for evaluating the organoleptic potentials of rice i.e., appearance, aroma, taste, and texture which are significant to consumer acceptance and marketability. This review consolidates current methodologies for sensory scoring of rice, including both human panel assessments and instrumental techniques (GC, Electronic nose, TPA, AI). Panelist training, standardized evaluation protocols, and the integration of sensory data into rice breeding and quality control programs may include. By examining global practices and the review underlines the sensory profiling in the variety approval process and highlights its role in developing rice cultivars that align with both agronomic performance and market expectations.

Keywords: Aroma, Basmati, sensory, KOH, Chromatography, texture.

INTRODUCTION

Rice (*Oryza sativa L.*) is an essential food for more than half the world's population. Beyond produce and nutritive content, organoleptic qualities such as flavor, aroma, and texture significantly impact consumer preference and marketability (Calingacion *et al.*, 2014). Sensory assessment delivers a systematic approach to measure subjective traits, enabling researchers and producers to improve varieties and processing methods. While agronomic traits such as yield and disease resistance remain crucial, sensory characteristics significantly influence market preference and adoption. National and international variety release committees increasingly emphasize sensory quality as part of rice variety approval. (Hossen *et al.*, 2021)

Sensory valuation plays a decisive role in rice breeding as it bridges the gap between agronomic performance and consumer preference. Whereas breeders conventionally focus on yield, pest resistance, and stress tolerance but the ultimate feat of

a rice variety in the market depends on appearance, aroma, texture, and taste (Bard., 2008; Bett-Garber *et al.*, 2017).

Consumers are cognizant to organoleptic traits, especially in premium varieties such as Basmati and Jasmine rice. Integrating personal perception into the breeding pipeline enables breeders to select lines that not only perform well in the field but also meet or exceed consumer expectations. (Juliano, 2016). Additionally, sensory profiling aids in the diversity of rice varieties for niche markets and confirms consistency as domestic and export standards. As international markets grow more competitive and consumer-driven, sensory evaluation has become a necessary factor of rice breeding programs. (Meullenet *et al.*, 2007; Champagne *et al.*, 2010; Tuano *et al.*, 2022).

Sensory evaluation of rice is based on four vital domains, each significantly influencing to consumer acceptance and marketability.

1. Appearance includes grain color, shape, chalkiness and translucency. These visual features influence consumer perception before cooking and can be a decisive reason in selection. (Unnevehr *et al.*, 1992). Recent studies confirm chalkiness and reduced translucency as key drivers of diminished market value and lower consumer preference (Xiao *et al.*, 2025).

2. Aroma is chiefly crucial for aromatic varieties like Basmati and Jasmine which is obsessed by volatile compounds especially 2-acetyl-1-pyrroline (2-AP). It refers to the scent of rice mutually before and after cooking. (Singh and Singh, 2000; Choudhury and

Article History

Received: [June 30, 2025](#), Accepted: [August 15, 2025](#),

Published: [September 06, 2025](#).




Copyright: © 2022 by the authors. Licensee Roots Press, Islamabad Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

<https://creativecommons.org/licenses/by/4.0/>

Gupta, 2014). Recent research reaffirms that 2-AP is the principal marker of aroma in fragrant cultivars, and that its biosynthesis is genetically regulated (e.g., via BADH2) and environmentally responsive during grain development and processing (Hashemi et al., 2025).



Texture comprises of mouthfeel, hardness, softness cohesiveness, stickiness and chewiness. It is typically evaluated after cooking and plays a key role in rice mouth feels and impacts overall satisfaction. (Tairi et al., 1992). A texture analyzer is an instrumental device used to objectively measure the physical properties of food, such as hardness, chewiness, and cohesiveness, providing precise and reproducible texture data. Recent studies have shown that higher adhesiveness (stickiness) correlates with increased amylopectin content, while higher hardness is associated with greater amylose levels (Li et al., 2023).

Taste/Flavor is the collective outcome of taste and aroma, resulting to the overall palatability of the rice. The equilibrium of umami, sweetness and minor bitterness impacts consumer's choice. (Bhattacharya and Corke, 1996; Tuano et al., 2022)

Sensory Evaluation Methods

Evaluation techniques are effective for defining the cooking quality and consumer adequacy of rice. It delivers a systematic process to quantify qualities such as texture, aroma, taste and appearance. Following techniques are commonly used in rice sensory appraisal system. (Hossen et al., 2021).

Descriptive Method

Trained panels assess predefined lexicon of rice flavor attributes i.e., nutty, floral or as quantitative and reproducible numerical results. It provides a comprehensive rice's sensory profile to enhance importance of aroma or texture. Feedback of descriptive analysis are numbered on predefined sensory scale for diverse rice varieties. (Stone and Sidel, 2004; Lawless and Heymann, 2010)

Discrimination methods/ Duo-Trio Test / Triangle Test

The Triangle Test comprise of three samples, two of which are the same and one different, and the trained participants are inquired to identify another sample. A reference sample is given in Duo-Trio Test and participants are allowed to select between two samples, one is the identical as the reference. These trials are suitable for detection of minor alterations in sensory traits i.e., aroma or texture of new varieties or standard. (ISO 2006, Luna 2018)

Texture Profile Analysis (TPA)

It is used to quantify texture qualities of rice i.e., cohesiveness, chewiness, elasticity and firmness. TPA is usually associated with equipment like a Texture Analyzer. It generates unbiased information that is correlated with the judgment of sensory panels. TPA provide valuable data on texture preferences that is tough to capture through human panels alone. Linking TPA and panel data consents an inclusive understanding of rice texture features. Wang et al. (2022) demonstrated significant correlations between instrumental hardness and stickiness with aroma compounds in cooked rice, while Choi and Seo (2023) reported that TPA-derived hardness, stickiness, and chewiness strongly influenced consumer liking across age and gender groups. Similarly, Janani et al. (2022) showed that parameters such as springiness and cohesiveness align with oral processing behaviors, and a post-2020 study on purple rice formulations confirmed that elasticity, hardness, chewiness, and cohesiveness significantly predict sensory evaluation scores.

Aroma Testing Using KOH (Potassium Hydroxide)

Potassium hydroxide (KOH) testing remains a widely used qualitative method for detecting aroma in aromatic rice varieties such as Basmati and Jasmine. The test primarily targets the presence of 2-acetyl-1-pyrroline (2AP), the key compound responsible for the characteristic popcorn-like scent of these rices. In this

method, rice grains or leaves are cooked or steamed to release volatile compounds, and a small amount of KOH solution—commonly 1.7%—is added to the sample or its vapor. The alkali enhances the volatility of 2AP, making the aroma more perceptible to trained evaluators. This approach, first described by Haug (1972) and later refined by Gupta et al. (2015), is favored in breeding programs for its simplicity, low cost, and rapid results, despite being subjective in nature. Recent research confirms that KOH testing is effective for rapid screening, though it is increasingly complemented by instrumental methods such as gas chromatography–mass spectrometry (GC–MS) for quantitative accuracy (Hien et al., 2021; Boontakham et al., 2022). Studies have shown that KOH-based aroma detection correlates well with sensory evaluation in distinguishing aromatic from non-aromatic genotypes, making it useful in early-generation selection (Roy et al., 2020). However, as environmental conditions and postharvest handling can influence 2AP concentration, advanced analytical techniques are often employed alongside KOH testing to provide a more robust understanding of aroma profiles. This integration ensures both rapid field-level screening and precise laboratory-based quantification, improving the efficiency of aromatic rice breeding and quality assessment.

Aroma profiling Using Gas Chromatography (GC)

Gas chromatography (GC), particularly when combined with head-space sampling techniques, is a widely used method for analyzing rice aroma because it enables the separation, identification, and quantification of volatile compounds, including trace levels of 2-acetyl-1-pyrroline (2AP) (Liu, 2013). In aromatic rice studies, GC methods such as static head-space GC coupled with flame ionization detection (GC-FID), nitrogen–phosphorus detection (GC-NPD), and mass spectrometry (GC-MS) have been successfully applied to generate detailed volatile profiles (Boontakham et al., 2022). These instrumental techniques provide high sensitivity, reproducibility, and specificity, making them indispensable for precise aroma characterization in quality control and breeding programs.

Despite its advantages, GC is constrained by several practical limitations. The equipment is costly, requires trained personnel for operation, and involves time-consuming procedures for sample preparation, chromatographic separation, and data analysis (Saxena et al., 2000; Jiang, 2008). Consequently, many resource-limited laboratories and small-scale industries face accessibility challenges, often relying on simpler qualitative assays such as the potassium hydroxide (KOH) test for preliminary screening (Roy et al., 2020). Integrating GC with rapid field-level methods offers a balanced approach—enabling both

large-scale screening and precise quantification of rice aroma compounds for research and commercial applications.

Advanced Techniques (Smart phone, electronic nose)

Smartphone-based aroma detectors and biosensors hold great potential to revolutionize rice quality testing by providing rapid, portable, and cost-effective tools for on-site aroma and flavor assessment. These devices leverage advances in sensor technology—such as electronic noses, gas sensors, and bio-recognition elements integrated with smartphones—to detect key volatile compounds for aroma only. Additionally, coupling biosensors with smartphone apps can facilitate data collection, analysis, and sharing, enabling large-scale sensory-instrumental correlations and consumer preference mapping. Although still emerging, these technologies promise to complement traditional sensory panels and instrumental methods, making rice quality evaluation more efficient, objective, and scalable (Banerjee et al., 2021; Li et al., 2023).

An electronic nose (e-nose) is an array of gas sensors designed to detect and differentiate complex volatile compounds, mimicking human olfactory perception, and is increasingly used for rapid aroma profiling in rice (Dutta et al., 2020)

Consumer Testing/untrained testing

Untrained panels or wide-ranging consumers are invited to judge the overall suitability of rice. Hedonic scaling or preference tests are employed. This process is mostly used for market research or real-world buyer preferences. (Bourne, 2002). Although it provides straight insights into how well a rice variety is likely to perform in the market but parameters such as taste, aroma, texture, and appearance are misled during overall experience. It is synergistically combined with other methods to refine rice varieties for broad consumer acceptance. (Champagne et al., 2010).

Data Collection, Analysis and Interpretation

Data composed from different panels are tabulated and analysed to quantify the overall strength of the aroma. Consumer preference for most rice varieties based on aroma. Principal component analysis (PCA) is used to quantify aroma preconceptions in different rice varieties. It supports the researchers and breeders to recognize how precise aroma traits are perceived by masses, which is decisive factor for evolving rice varieties that appeal to specific markets. (Lestari et al., 2011)

Composition of the Panel and training

Sensory testing of rice includes diverse panelists, each examining specific quality trait based on their proficiency. A proficient panel typically entails 5-10 persons (knowing hedonic protocols of rice). A semi-trained panel contains 10- 15 members (having some

understanding in examining rice taste) frequently guided on scoring to enhance reliability. (Meilgaard et al., 2015)

Whereas a buyer panel / un trained contains 30 - 50 individuals from the general population and is mostly used in acceptability or liking/ disliking of varieties to assess how this specific variety will perceived by common masses. Nominated participants/panel must be unbiased, impartial, non-smokers and healthy to sustain 982586aroma sensitivity. (Delwiche, 2004)

Real estimation depends on the skill of a proficient sensory panel. Appropriate training confirms that the participants are able to assess rice cooking parameter’s accurately and precisely. Panelists must acquaint with reference standards that demonstrate attributes i.e., nutty, floral or sticky. Furthermore, it is vital for all participants to develop harmony to avoid variations. Strong understanding of parameters i.e., chewy, nutty, flowery, soft, hard, aromatic or firm confirms interpretation. Calibration using identified samples with pre-defined proforma is necessary to perceive attributes accurately. (Lawless 2010; ASTM 2011).

Training includes repeated exposure to standard samples, calibration exercises, and consensus discussions to minimize individual variability. Cross-validation involves periodically testing panelists' evaluations against each other and against instrumental measurements or expert benchmarks to assess consistency, reproducibility, and alignment with objective criteria. (Stone & Sidel, 2004)

Standardized Sample Preparation Procedure

25g uncooked rice samples are used per sample per replicate. Rice samples are soaked in distilled water

for 30 minutes prior cooking. Cooking is carried out using 1:2 rice-to-water ratio in boiling water bath. After cooking, the rice samples are allowed to cool for 5-10 minutes before evaluation. To confirm unbiased assessment, all samples are blind-coded using random numbers for secrecy. Panelists estimate all parameters independently and record scores using a hedonic scale standardized for Basmati/ coarse varieties. For holistic understanding of rice quality traits both human perception and equipments are combined to ensure accuracy and precision, although sensory evaluation merely insights into customer acceptability. (Moskowitz and Gofman 2006).

Lexicon development for sensory descriptors in Basmati rice involves creating a precise, standardized vocabulary to describe its unique quality traits, including grain appearance (slender shape, whiteness, translucency), characteristic popcorn-like aroma from 2-acetyl-1-pyrroline, delicate nutty or floral notes, and cooked grain texture attributes such as softness, fluffiness, and non-stickiness. This process requires selecting representative Basmati samples, engaging trained sensory panels to generate and refine terms, defining each descriptor with clear definitions and reference standards, and establishing intensity scales to ensure consistency across evaluations. Such a lexicon enables breeders, quality controllers, and exporters to communicate quality objectively, link sensory traits with instrumental measurements, and maintain Basmati’s premium identity in domestic and global markets (Cruz & Khush, 2000; IRRI, 2024; Matsuda et al., 2024).

Sensory Descriptor	Attribute	Details
Appearance	Slender Grain	Long, thin grain shape / Basmati
	Chalkiness	Opaque white patch in the grain due to loosely packed starch granules.
Aroma	Popcorn-like	Distinct nutty-popcorn aroma due to 2-acetyl-1-pyrroline (2-AP)
	Nutty	Aroma reminiscent of roasted nuts.
	Floral	Light, fragrant aroma similar to jasmine
	Sweet Aromatic	Mild sugary scent perceived in freshly cooked Basmati.
Flavor	Nutty Flavor	Roasted nut taste after chewing.
	Sweetness	Natural sweetness in taste after chewing cooked grains.
	Starchy Flavor	Gelatinized starch taste typical of cooked rice.
Texture (Cooked)	Fluffiness	Individual grains remain separate, light, and airy after cooking
	Softness	Low resistance to compression during chewing.
	Non-stickiness	Lack of grain adhesion after cooking.
	Firmness	Resistance to deformation on chewing without hardness.

Future Directions

Sensory evaluation of rice grains plays a critical role in the modern food science sectors. It allows breeders, food scientists and industry stakeholders by scientifically assessing the organoleptic qualities to rise the competitiveness of rice in local and international markets. As global markets become more diverse, the combination of human perception and instrumental methods will expand the accuracy and rice evaluation. Sensory evaluation should be included as mandatory portion of rice breeding program.

Future research should focus on establishing 2-3 training programs for sensory panels and cross-cultural research to ensure international pertinency. In conclusion, sensory evaluation not only assists as a scientific tool for refining rice quality but also as a bridge between field research and consumer satisfaction, ensuring the sustainable achievement of rice in a competitive global market.

Conflict of Interest

The authors declare no conflict of interest

REFERENCES

- American Society for Testing and Materials (ASTM). (2011). *ASTM E1958-11: Standard guide for sensory evaluation training*. ASTM International.
- Banerjee, S., Das, A., & Mandal, P. (2021). Smartphone-based sensors for food quality assessment: A review. *Trends in Food Sci & Technol*, 112, 69–79. <https://doi.org/10.1016/j.tifs.2021.02.005>
- Bard, L. E., & Gough, A. J. (2008). Sensory analysis of Basmati rice: A review of methods for evaluating aroma characteristics. *Food Quality and Pref*, 19(7), 554-564.
- Bett-Garber, K.L., Bryant, R.J., Grimm, C.C., Chen, M., Lea, J.M., McClung, A.M. (2017). physicochemical and sensory analysis of US rice varieties developed for the basmati and jasmine markets. *Cereal Chem.* 94(3), 602-610.
- Bhattacharya, K. R., & Corke, H. (1996). Sensory analysis of cooked rice. *Food Quality and Preference*, 7(3), 201–207.
- Boontakham, P., Chaiwan, P., & Sirisena, S. (2022). Advances in analytical techniques for aromatic rice aroma profiling. *Food Chem*, 389, 133057.
- Bourne, M. (2002). *Food texture and viscosity: Concept and measurement* (2nd ed.). Food Science & Technology, International series, Academic Press, Elsevier, USA. PP: 369
- Calingacion, M., Laborte, A., Nelson, A., Resurreccion, A., Concepcion, J. C., Daygon, V. D., Mumm, R., & Fitzgerald, M. A. 2014. Diversity of global rice markets and the science required for consumer-targeted rice breeding. *PLOS ONE*, 9(1), 1-12.
- Champagne, E. T., Bett-Garber, K. L., Fitzgerald, M. A., Grimm, C. C., Lea, J., & Ohtsubo, K. (2010). Important sensory properties differentiating premium rice varieties. *Rice*, 3(4), 270–281.
- Choi, S., & Seo, H. (2023). Influence of rice textural properties on consumer acceptance across age and gender. *Food Quality and Preference*, 105, 104887. <https://doi.org/10.1016/j.foodqual.2022.104887>
- Choudhury, R., & Gupta, R. K. (2014). Aroma compounds of Basmati rice: Chemistry and analysis. *Critical Reviews in Food Science and Nutrition*, 54(10), 1413-1422.
- Cruz, N. D., & Khush, G. S. (2000). Rice grain quality evaluation procedures. In R. K. Singh, U. S. Singh, & G. S. Khush (Eds.), *Aromatic rices* (pp. 15–28). New Delhi: Oxford & IBH Publishing.
- Delwiche, J. F. (2004). The impact of perceptual interactions on perceived flavor. *Food Quality and Preference*, 15(2), 137–146.
- Dutta, S., Sharma, H., & Kumar, S. (2020). Electronic nose applications for food aroma evaluation: A review. *Trends in Food Sci & Technol*, 97, 234–247.
- Gupta, M., & Verma, R. K. (2015). *Biotechnology and Aroma in Rice* (pp. 67-79). In M. R. Cereal Science and Biotechnology. Springer.
- Hashemi, M., Zhang, L., & Kumar, A. (2025). Genetic and environmental regulation of 2-acetyl-1-pyrroline biosynthesis in aromatic rice. *J of Cereal Sci*, 115, 104987.
- Haug, W., & L. L. L. (1972). A method for the evaluation of rice aroma. *Cereal Chem*, 49(5), 541-549.
- Hien, N. T., Thao, N. P., & Liem, T. H. (2021). Evaluation of KOH test and GC-MS for rapid detection of aromatic rice. *J of Cereal Sci*, 102, 103339.
- Hossen, M. J., Sreenivasulu, N., & Henry, R. 2021. Review of sensory evaluation and consumer acceptance studies of rice. *Cereal Chem.*, 98(5), 976–990.
- International Organization for Standardization (ISO). (2006). *ISO 4120:2006 - Sensory analysis: Triangle test*. International Organization for Standardization.
- International Rice Research Institute (IRRI). (2022). *Standard Evaluation System (SES) for Rice* (6th ed.).

- Janani, A., Kumar, S., & Ramaswamy, H. (2022). Oral processing behavior and its relation to rice texture parameters. *LWT – Food Sci & Technol*, 153, 112520.
- Jiang, J., & Xu, L. (2008). Aroma and volatile compounds in rice: new insights into the role of volatile compounds in rice aroma using gas chromatography-mass spectrometry (GC-MS). *J Food Sci*, 73(9): S472-S479.
- Juliano, B. O. 2016. *Encyclopedia of Food Grains* (2nd ed.). Philippine Rice Research Institute LosBañõs, Laguna, Philippines.1: 125-129. Oxford: Academic Press
- Lawless, H. T., & Heymann, H. (2010). *Sensory evaluation of food: Principles and practices* (2nd ed.). Springer Science & Business Media pp:81.
- Lestari, P., Singh, R. K., & Subudhi, P. K. (2011). Quantitative trait loci analysis for grain aroma in rice using a recombinant inbred population. *Mol. Breeding*, 27(4): 503–512.
- Li, Y., Zhang, Q., & Wang, J. (2023). Advances in biosensors for food aroma detection: Applications and challenges. *Biosensors and Bioelectronics*, 215, 114537.
- Liu, H., Chen, C., & Ren, J. (2013). Extraction and quantification of volatile aroma compounds from rice using solid-phase microextraction (SPME) and gas chromatography-mass spectrometry (GC-MS). *Food Res Int*, 51(1): 31-38.
- Luna, R. L., & R. H.M. (2018). Application of texture profile analysis (TPA) in the evaluation of cooked rice texture. *Food Res Int*, 105: 569-575
- Matsuda, T., Yamamoto, M., Sakamoto, T., & Ohtsubo, K. (2024). Development of a sensory lexicon for cooked rice. *J of the Japanese Society for Food Sci & Technol*, 72(6), 215–225.
- Meilgaard, M., Civille, G. V., & Carr, B. T. (2015). *Sensory evaluation techniques* (5th ed.). CRC Press.
- Meullenet, J. F., Xiong, R., & Findlay, C. J. 2009. Multivariate and Probabilistic Analyses of Sensory Science Problems. *Trends in Food Sci. Tech.* 20(2): 108-109.
- Moskowitz, H. R., & Gofman, A. (2006). Global sensory evaluation: Cultural differences and implications for product development. *Food Quality and Pref.*, 17(6): 531–539.
- Roy, S., Das, A., & Singh, V. (2020). Correlation between KOH-based screening and sensory evaluation for aromatic rice genotypes. *Int J of Food Sci & Technol*, 55(5), 1783–1792.
- Saxena, P. R., & Srivastava, M. (2000). Quantitative analysis of rice aroma in Basmati rice using gas chromatography-mass spectrometry (GC-MS). *Food Chem.* 71(2), 225-229.
- Singh, R. K., & Singh, U. S. (2000). *Aromatic Rices*. Oxford & IBH Publishing.
- Stone, H., & Sidel, J. L. (2004). *Sensory evaluation practices* (3rd ed.). Elsevier Academic Press. pp: 99-104
- Taira, H., Fujii, K., & Ishikawa, M. (1992). Texture and eating quality of cooked rice in relation to water content and amylose content. *J of Texture Studies*, 23(1), 55–68.
- Tuaño, A. P. P., et al. (2022). Electronic nose as a tool for discriminating aromatic rice varieties. *Food Control*, 136, 108878.
- Unnevehr, L. J., Duff, B., & Juliano, B. O. 1992. Consumer demand for rice grain quality: Terminology, measurement, and evidence. In B. O. Juliano (Ed.), *Rice chemistry and technology* (pp. 297–320). LosBañõs, Laguna, Philippines
- Wang, X., Li, Q., & Zhang, H. (2022). Correlation between rice textural properties and aroma compounds. *J of Food Sci*, 87(5), 1784–1794. <https://doi.org/10.1111/1750-3841.16123>
- Xiao, C., Zhang, H., Liu, F., & Sun, X. (2025). Sensory quality analysis of rice based on fuzzy sensory evaluation. *Food Sci and Technol*, 45(2), 123–134. <https://doi.org/10.13386/j.issn1002-0306.2017.21.049>